

gadolinium



Stable	Atomic mass*	Mole
isotope		fraction
¹⁵² Gd	151.919 791	0.0020
154 Gd	153.920 8656	0.0218
155 Gd	154.922 622	0.1480
156 Gd	155.922 1227	0.2047
157 Gd	156.923 9601	0.1565
158 Gd	157.924 1039	0.2484
¹⁶⁰ Gd	159.927 0541	0.2186

^{*} Atomic mass given in unified atomic mass units, u.

Half-life of redioactive isotope
Less than 1 second



Important applications of stable and/or radioactive isotopes

Isotopes in medicine

- 1) Gadolinium has the highest neutron cross-section among any stable nuclides, 61,000 barns for ¹⁵⁵Gd and 259,000 barns for ¹⁵⁷Gd. ¹⁵⁷Gd has been used to target tumors in neutron therapy. This element is very effective for use with neutron radiography and in shielding of nuclear reactors.
- 2) ¹⁵³Gadolinium is produced in a nuclear reactor from elemental europium or enriched gadolinium targets. It has a half-life of 240±10 days and emits gamma radiation with strong peaks at 41 keV and 102 keV. It is used in many quality assurance applications, such as line sources and calibration phantoms, to ensure that nuclear medicine imaging

systems operate correctly and produce useful images of radioisotope distribution inside the patient. It is also used as a gamma ray source in X-ray absorption measurements or in bone density gauges for osteoporosis screening, as well as in the Lixiscope portable X-ray imaging system.




Figure 1: This is a picture of a patient undergoing neutron therapy. The red lasers cross to target the patients' tumor. A high beam of neutrons is fired at the target to stop the growth and eradicate the tumor.

^{**}Applications of gadolinium isotopes are still being researched and this page will be updated shortly. **